



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

specimens from the following collections: K. Pr. Geol. Landesanstalt, Berlin; Städt. Museum, Chemnitz; Naturw. Museum, Dortmund; K. K. Geol. Reichsanstalt, Wien; Märkisches Museum, Witten; Museum, Osnabrück; s'Rijks Herbarium, Leiden; Geol. Institut der Universität, Göttingen; Geol. Institut der Bergakademie, Clausthal; and Geol. Institut der Universität, Münster i.W.

The work is strictly taxonomic. The descriptions and synonymy are very complete, and the magnificent plates enable one to study the subject almost as well as if he had the actual specimens in his hands.—CHARLES J. CHAMBERLAIN.

Anatomy of petioles of Cycads.—LE GOC¹⁷ has studied the debated xylem situation in the foliar bundles of the cycads. As is well known, both centripetal and centrifugal xylem occur, a situation which has been variously interpreted, but in general the bundle has been spoken of as mesarch. LE GOC has found that at the very base of the petioles in cycads the xylem is entirely centrifugal, and that later the bundles assume different forms, becoming concentric, collateral, or a combination of these two arrangements. He regards the centrifugal xylem at the base as a secondary growth, and the centripetal xylem which appears later as a primary structure, laid down early, but only gradually lignified. The two xylems he thinks are probably distinct in origin, appearing in response to "physiological demands but morphologically discontinuous." During most of the course of the bundles along the petiole the two xylems remain distinct, and therefore he would suggest that the bundle is more properly called "pseudo-mesarch."—J. M. C.

The origin of the "eye spot."—From a study of the literature, without any apparent study of the structures themselves, ROTHERT¹⁸ comes to the conclusion that the "eye spot" of flagellates and algae is derived from a plastid. The evidence seemed clear already, so far as the Euglenaceae are concerned, and GUIGNARD, as long ago as 1889, claimed that the "eye spot" seen in the sperms of *Fucus* arose from a colorless plastid. ROTHERT overlooks entirely the various papers by YAMANOUCHI from 1909 to 1913, particularly the paper on *Culleria*,¹⁹ in which the origin and development of the "eye spot" are described in detail.—C. J. CHAMBERLAIN.

¹⁷ LE GOC, M. J., Observations on the centripetal and centrifugal xylems in the petioles of cycads. *Ann. Botany* 28:183-193. *pl. II. fig. I.* 1914.

¹⁸ ROTHERT, WLADISLAW, Der "Augenfleck" der Algen und Flagellaten ein Chromoplast. *Ber Deutsch. Bot. Gesells.* 32:91-96. 1914.

¹⁹ YAMANOUCHI, SHIGÉO, The life history of *Culleria*. *BOT. GAZ.* 54:441-502. *figs. 16. pls. 26-45.* 1912.